Patient name:
DOB:
NHS number:

- Carers must be given training in administration of midazolam. Midazolam may not be prescribed for use by carers in some areas due to local policy.
- Please ensure, with the help of the GP and specialist team, that medications and dosages are kept up to date.

	For carer use		For healthcare professional use						
Medication	**Midazolam** Brand:	**Lorazepam** Brand:	**Morphine sulphate** Brand:	**Glycopyrronium bromide** Brand:	**Hyoscine hydrobromide** Brand:	**Midazolam** Brand:	**Cyclizine** Brand:	**Haloperidol** Brand:	**Levomeprom-azine** Brand:
Used for	• restlessness • agitation • fear • anxiety	• restlessness • agitation • fear • anxiety	• pain • breathlessness	• excessive/noisy respiratory secretions • excessive saliva	• excessive/noisy respiratory secretions • excessive saliva	• restlessness • agitation • fear • anxiety	• nausea	• nausea • agitation • restlessness	• nausea • restlessness
Administration	Buccal (into mouth between cheek and gums)	Tablet or sublingual (under the tongue)	Subcutaneous injection **or** infusion	Subcutaneous injection **or** infusion	Transdermal patch **or** infusion	Subcutaneous injection **or** infusion	Subcutaneous injection **or** infusion **or** tablets	Tablets **or** capsules **or** subcutaneous injection **or** infusion	Tablets **or** subcutaneous injection **or** infusion
Tick if included in this box.	☐ Dosage:	☐ Dosage:	☐ Dosage:	☐ Dosage:	☐ Dosage:	☐ Dosage:	☐ Dosage:	☐ Dosage:	☐ Dosage: